Imagerie cérébrale et Langage

Introduction : Langage et cerveau

Aphasies et langage

Hémisphère GAUCHE
Gyrus frontal inférieur (3eme
circonvolution frontale; aires
44 et 45)

Hémisphère GAUCHE
partie postérieure du gyrus temporal supérieur
(cortex auditif « associatif », aire 22 de
Brodmann)

Figure 20.2 bear, faisceau arqué.
Cheminement de l'information pour la
répétition d'un mot

Modèle de Wernicke Geschwind

B- Localisation du langage

Quelques problèmes avec le modèle de Lichtheim

- Les aphasiques de Broca ont des difficultés de compréhension et inversement
- La sévérité des aphasies de Broca et Wernicke dépend de l'étendue de la lésion
- L'aphasie est influencée par les dommages dans les structures sous-corticales (par exemple thalamus)
- Les cas d'aphasie après excision d'une tumeur ou d'une zone épileptique sont en général plus légers que les aphasies après AVC, qui impliquent des régions plus étendues
- Il y a souvent récupération après un AVC car d'autres parties du cerveau non atteintes peuvent compenser

Stimulation cérébrale

(A)

(B)

FIGURE 10-19 Mapping Cortical

Functions. (A) The patient is fully conscious during neurosurgery for intractable epilepsy,

Stimulation du cortex moteur: arrêts de la parole ou parfois vocalisations

Stimulations d'autres parties du cerveau pendant la parole: arrêts, hésitations, confusions, difficultés à trouver le mot

Stimulations électriques du cortex cérébral pour la cartographie des aires du langage au cours de la dénomination d'images

Penfield et Roberts, 1959

Ojemann, 1989
(< 100 patients)

- * Variabilité des localisations
- * Beaucoup de régions extérieures aux « aires du langage »

; dans Neurosciences, Purves et al.

Imagerie cérébrale

Premières études en TEP

Petersen et coll, 1988

Figure 1. Four different hierarchically organized conditions are represented in these mean blood flow difference images obtained with PET. All of the changes shown in these images represent increases over the control state for each task. A group of normal subjects performed these tasks involving common English nouns (Petersen et al. 1988; Petersen et al. 1989) to demonstrate the spatially distributed nature of the processing by task elements going

complexity was increased from simply opening the eyes (row 1) through passive viewing of nouns on the television monitor (row 2); reading aloud the nouns as they appear on the screen (row 3); and saying aloud an appropriate verb for each noun as it appeared on the screen (row 4). These horizontal images are oriented with the front of the brain on top and the left side to the reader's left. The marking "Z = 40" indicates millimeters above and below a horizontal plane

I Caractéristiques des méthodes d'imagerie cérébrale

1- Structurale / Fonctionnelle

2- Temps / espace

- Le *pouvoir de résolution* est la plus faible variation de la quantité mesurée que l'instrument soit capable de mettre en évidence

Figure 3.7 The cell membrane: A lipid bilayer with signal proteins and channel proteins embedded in it.

Espace: de moins d'un nM à plusieurs dizaines de cm

Temps: de quelques μ s à plusieurs années

II- Imagerie structurale

Principes de la résonance magnétique nucléaire

(Cours de M. Roth)

http://irmfmrs.free.fr/formation/cours_irm/cours_irm.htm

- L'IRM exploite les propriétés magnétiques des noyaux atomiques

Imagerie en tenseur de diffusion

distribution des directions de diffusion des molécules d'eau. Cette diffusion étant contrainte par les tissus environnants, cette modalité d'imagerie permet d'obtenir indirectement la position, l'orientation et l'anisotropie des structures fibreuses,

III Méthodes d'imagerie FONCTIONNELLE

1 Méthodes de mesure du débit sanguin cérébral

TEP et IRM fonctionnelle

Fin 19eme siècle: Mosso

1928: Fulton

Le signal BOLD

Hémoglobine ou Désoxyhémoglobine

(A)

Pure oxygen

(B)

Normal Air

L'augmentation du taux d'hémoglobine déoxygénée fait chuter le signal (Ogawa et al., 1990a chez le rat)

La déoxyhémoglobine est paramagnétique (Pauling 1936): elle crée des inhomogénéités locales du champ magnétique → influence sur les paramètres mesurés en RMN

Kwong et collaborateurs, 1992

Humain, stimulation visuelle

Décalage de la réponse de quelques secondes après le début/la fin de la stimulation

Le paradigme d'activation

- Mesurer la pensée: Donders 1868

180 ms

syllabe
(toujours la
même)

Réponse

268 ms

syllabe
(variable)

Réponse

Etape d'identification : 88 ms

Insertion pure

→ le paradigme d'activation

- Le débit sanguin local ne varie pas suffisamment à cause d'une opération mentale donnée pour pouvoir être discriminé de l'activité globale du cerveau
- La mesure dans la condition d'intérêt est toujours considérée relativement à la mesure dans une autre condition

Etudes de groupe

IRMf : méthode indirecte

- Le « filtre hémodynamique » : **Réflexion *indirecte***
- **de l'activité neuronale**

- Couplage réponse BOLD/ activité neuronale:
- relation complexe → Variations? Âge, pathologie,
- pharmacologie

2- Méthodes de mesure de l'activité électrique corticale

L'électroencéphalographie

EEG

- Berger, 1929, chez l'homme (sur un patient avec un
- défaut du crâne)
- Mesure de l'activité électrique du cortex cérébral
- Nombre variable d'électrodes (16 à 128) couvrant
- l'ensemble du scalp (F, C, T, P, O) sur emplacements standards

- Différences de potentiel enregistrées entre deux
- électrodes sur le scalp
- Amplificateurs différentiels
- DDPs = quelques microvolts
- (μV)

- Chaque ampli enregistre la
- différence entre une électrode
- et l'électrode dite « de référence »

- Méthode non invasive
- Nomenclature des électrodes

- Dérivation bipolaire

- Dérivation avec référence commune (non sensible aux variations de potentiels cérébraux),

Synchronisation de l'activité des neurones générant un EEG de forte amplitude

Décharge irrégulière

(b)

Décharge synchronisée

(c)

C1: le mode spontané

- signal continu

- Observation des rythmes cérébraux (oscillations)

C2- Potentiels évoqués

- Technique de moyennage:
- temps 0 = évènement d'intérêt
- Cet évènement est répété plusieurs fois
- Les signaux enregistrés à chaque répétition sont moyennés

Chaque courbe montre l'activité électrique moyenne enregistrée pendant une seconde après la présentation du mot visuel, mais le nombre d'essais utilisés pour calculer le moyennage augmente de haut en bas. Avec un seul essai, le bruit est très élevé, mais il est fortement réduit à mesure qu'un plus grand

Kutas et Hillyard, 1980

Applications au langage

La lecture

Figure 7. Correlation between the lesion site associated with pure alexia (top) and the left occipito-temporal activation during normal reading (bottom). The normalized lesions from six patients were combined, to compute for each voxel an index proportional to the log likelihood ratio, expressing the conditional link between the existence of a lesion and the occurrence of pure alexia. The critical area follows the left occipito-temporal sulcus, and overlaps neatly with the VWFA, as identified with fMRI in normal subjects by contrasting alphabetic stimuli versus checkerboards.

Alphabetic stimuli vs fixation

Alphabetic vs fixation

Alphabetic vs chequerboards

Words vs consonants

Consonants vs words

Stimulated hemifield

Fig. 7 Topography of the subtracted ERPs to words and non-words, and average potentials recorded on a left temporal electrode (white square) in control subjects and patient A.C. ANOVAs were performed on voltages averaged over a 240–360 ms time-window (rectangles). In controls word stimuli (green curves) elicited a prolonged left temporal negativity relative to non-words (red curves), irrespective of the stimulated hemifield, but only following RVF stimulation in patient A.C.

How Learning to Read Changes the Cortical Networks for Vision and Language

Stanislas Dehaene, *et al.*
Science **330**, 1359 (2010);
 DOI: 10.1126/science.1194140

VWFA
 Response at [-44, -50, -14]

Left frontal
 Response at [-46, -2, 52]

Occipital
 Response at [22, -86, -10]

Response at [-54, 26, -6]

Left superior temporal sulcus

Response at [-50, -44, 6]

Response at [-50, 12, -24]

Sujets lettrés, illettrés,
 ex-illettrés portugais ou
 brésiliens

A Modulated activation of *Planum Temporale* to spoken language

Spoken sentences in localizer

Response at [-38, -28, 18]

Auditory lexical decision

Response at [-38, -28, 18]

Figure 3. The organization of visual areas differs in good readers and in dyslexics (redrawn from Monzalvo et al., 2012). Good readers show a well-developed visual word form area (shown in green), a region that responds to written words more than to other categories of visual stimuli (faces, places, or checkerboards). Dyslexics show no such specialization for written words, and also exhibit a much weaker activation to faces in the right hemisphere. The evidence suggests that literacy involves a specialization of the visual word form area in the left hemisphere and, as a result, the displacement of face responses to the right hemisphere—two processes that fail to occur in young dyslexics. - See more at: http://dana.org/Cerebrum/2013/Inside_the_Letterbox__How_Literacy_Transforms_the_Human_Brain/#sthash.ENcXEaBE.dpuf

Implicated in dyslexia: The arcuate fasciculus is an arch-shaped bundle of fibers that connects the frontal language areas of the brain to the areas in the temporal lobe that are important for language (left). Researchers found that kindergarten children with strong pre-reading scores have a bigger, more robust and well-organized arcuate fasciculus (bottom right) while children with very low scores have a small and not particularly well-organized arcuate fasciculus (top right). (Zeynep Saygin/MIT)

Origines et latéralisation du langage

Meta-analyses

Vigneau et al. 2003

Price, 2012

Figure I. Examples of anatomical factors explaining part of the inter-individual variability of the left HS for language. (a) Correlation between the functional hemispheric asymmetry index and head size indexed by skull perimeter during story listening compared to a story in a language unknown to the participants. The larger the brain of the right-handed participants, the stronger the leftward asymmetry. Reproduced, with permission, from [85]. (b) Familial sinistrality (FS) has a significant impact on the left planum temporale (LPT) surface area. Subjects with a left-handed first degree relative (FS+) have a smaller LPT surface area than subjects who do not, independent of their handedness. Note that subjects with left-handed mothers have the smallest LPT surface area. Reproduced, with permission, from [83]. (c) Correlation between gray-matter density in Heschl's gyrus and 2-Hz frequency modulation detection threshold at age 10 overlaid on a T1-weighted average image template of participants at age 11.5. Colored areas represent the value of the *t* statistic for voxels where gray-matter density significantly correlates with auditory FM threshold score; the peak in the left Heschl's gyrus is marked with a cross. Reproduced, with permission, from [88]. (d) Left: group average covariation map during temporal processing of sounds; middle: correlation between left Heschl's gyrus volume and the extent of temporal-related activity within left Heschl gyrus; right: absence of correlation between activation and right Heschl's gyrus volume. Reproduced, with permission, from [86].

a. Whole-brain analyses of inter-hemispheric asymmetries

b. Inter-hemispheric asymmetries of the superior temporal sulcus

Exemple: le langage chez les bébés

- Dehaene-Lambertz (1994)

Les potentiels évoqués (ERP) des enfants autistes qui ont une préférence pour les sons n'appartenant pas au langage sont altérés

Absence de MMN (*mismatch negativity* – négativité de discordance) chez les enfants autistes préférant des sons qui ne sont pas issus du langage (enregistrements dans le lobe temporal)

x

[Species-specific calls activate homologs of Broca's and Wernicke's areas in the macaque](#)

Ricardo Gil-da-Costa, Alex Martin, Marco A Lopes, Monica Muñoz, Jonathan B Fritz & Allen R Braun
Nature Neuroscience 9, 1064 - 1070 (2006)